

University of Groningen

Unprecedented catalytic enantioselective trapping of arene oxides with dialkylzinc reagents

Bertozzi, F.; Crotti, P.; Del Moro, F.; Feringa, B.L.; Macchia, F.; Pineschi, M.

Published in:
Chemical Communications

DOI:
[10.1039/b108541g](https://doi.org/10.1039/b108541g)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2001

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Bertozzi, F., Crotti, P., Del Moro, F., Feringa, B. L., Macchia, F., & Pineschi, M. (2001). Unprecedented catalytic enantioselective trapping of arene oxides with dialkylzinc reagents. *Chemical Communications*, 113(24), 2606 - 2607. <https://doi.org/10.1039/b108541g>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Supplementary Material
for
Unprecedented Catalytic Enantioselective Trapping of Arene Oxides
with Dialkylzinc Reagents.

Fabio Bertozzi, Paolo Crotti, Federica Del Moro, Franco Macchia, Mauro Pineschi*
Dipartimento di Chimica Bioorganica e Biofarmacia, Università di Pisa,
Via Bonanno 33, 56126 Pisa, Italy

Ben L. Feringa*
Department of Organic and Molecular Inorganic Chemistry, Stratingh Institute,
University of Groningen, Nijenborgh 4, NL9747 AG Groningen, The Netherlands.

General. All reactions were conducted in flame dried glassware with magnetic stirring under an atmosphere of argon. Toluene was distilled from sodium/benzophenone ketyl and stored under argon. Et₂Zn (1.1 M solution in toluene) and Me₂Zn (2.0 M solution in toluene) were purchased from Aldrich. Analytical TLC were performed on Alugram SIL G/UV254 silica gel sheets (Macherey-Nagel) with detection by 0.5% phosphomolybdic acid solution in 95% EtOH. Silica gel 60 (Macherey-Nagel 230-400 mesh) was used for flash chromatography. Solvents for extraction and chromatography were HPLC grade.

Optical rotation were measured on a Perkin-Elmer 241 digital polarimeter with a 1 dm cell. ¹H NMR spectra were recorded on a Bruker AC-200 spectrometer on CDCl₃ solution. Chemical shifts are reported in ppm downfield from tetramethylsilane with the solvent resonance as the internal standard (deuteriochloroform: δ 7.26). ¹³C NMR spectra were recorded on a Bruker AC-200 (50 MHz) spectrometer with complete proton decoupling. Chemical shifts are reported in ppm downfield from tetramethylsilane with the solvent resonance as the internal standard (deuteriochloroform: δ 77.7). Gas chromatography was performed on a Perkin-Elmer 8420 apparatus (FI detector) using a Chromopak fused silica 25 m x 0.25 mm column, coated with CP-Cyclodextrin-B-236-M-19). In all cases, the injector and detector temperature was 250°C and a 1.8 mL / min helium flow was employed. Analytical high performance liquid chromatography (HPLC) was performed on a Waters 600E equipped with a Waters 990 photodiode array detector using a Daicel Chiralcel OD-H column.

A typical procedure for the enantioselective trapping of arene oxides with dialkylzinc reagents. Formation of **3a (**5a**).**

A 25 ml Schlenk tube under argon was charged with Cu(OTf)₂ (5.8 mg, 0.015 mmol), (*R,R,R*)-**2** (16.2 mg, 0.03 mmol), anhydrous toluene (2 ml), and was stirred at room temperature for 40 min. The colorless solution was cooled to -78°C and subsequently additioned with a solution of arene oxide **1a** (94 mg, 1.0 mmol) in toluene (0.5 ml) and Me₂Zn (0.75 ml). The reaction was followed by GC analysis and quenched with saturated aqueous NH₄Cl after 1h (95% conversion). Extraction with Et₂O and evaporation of the dried (Na₂CO₃) organic phase gave a crude product (106 mg) essentially constituted by the two regioisomers **3a** and **5a** in a 69:31 ratio and the chiral ligand (*R,R,R*)-**2**. In our hands, it was not possible to isolate in a pure state the achiral γ -adduct **5a**. However the presence of **5a** as a reaction product was clearly indicated by ¹H and ¹³C NMR of the mixture. **5a**: ¹H NMR δ (CDCl₃): 5.97-5.7 (m, 4H), 4.49-4.37 (m, 1H), 2.72-2.56 (m, 1H), 0.98 (d, 3H, *J* = 7.3 Hz). ¹³C NMR δ (CDCl₃): 127.7, 122.3, 62.6, 38, 21.9. After repeated chromatographic purification (SiO₂, hexanes/Et₂O=85:15 plus 1% NEt₃) it was possible to obtain an analytical pure sample of **3a** (8 mg).

(1*S*, 6*S*)-6-Methyl-2,4-cyclohexadien-1-ol (3a): a colourless liquid. ¹H NMR δ (CDCl₃): 5.97-5.70 (m, 4H), 3.97-3.87 (m, 1H), 2.52-2.33 (m, 1H), 0.96 (d, 3H, *J* = 7.3 Hz). ¹³C NMR δ (CDCl₃): 134.2, 133.4, 127.2, 125.4, 71.4, 31.5, 17.8.

The enantiomeric excess of **3a** (93%) was determined by chiral GC (CP-cyclodex- β -column), isothermal 107 °C, (1*S*, 6*R*) (minor) t_R14.2 min, (1*R*, 6*S*) (major) t_R 14.7 min.

Absolute configuration of **3a** was determined by comparison of optical rotation after conversion (H₂/Pd/AcOEt on charcoal) into *trans*-2-methylcyclohexanol, a compound of known absolute configuration.¹

¹ M.P. Doyle, A.V. Kalinin, D.G. Ene *J. Am. Chem. Soc.* 1996, **118**, 8837.

(1S, 6S)-(+)-6-Methyl-2,4-cyclohexadien-1-benzoate (3a-Bz): to a solution of a 64:36 mixture of **3a** and **5a** (40 mg, 0.36 mmol) in CH₂Cl₂ (2ml) and pyridine (0.5 ml), benzoyl chloride (41 μ l, 0.65 mmol) was added at 0°C under stirring. The mixture was kept at 0°C for 2 h, allowed stand at room temperature for 24h, and then quenched at 0°C with H₂O maintaining the stirring for further 2h. The solution was then diluted with CH₂Cl₂, washed with water, NH₄Cl saturated solution, NaCl saturated solution and dried over MgSO₄. Only benzoylated **3a** was found as a product and the crude reaction mixture was purified by flash chromatography (hexanes: AcOEt=85:15) to deliver **3a-Bz** (39 mg, 80% yield). $[\alpha]_D^{23}=+267.3$ ($c=2.75$, CH₂Cl₂), ¹HNMR (CDCl₃): 8.08-8.01 (m, 2H), 7.58-7.38 (m, 3H), 6.18-6.11 (m, 1H), 6.03-5.84 (m, 3H), 5.43 (t, 1H, $J = 5.1$ Hz), 2.81-2.64 (m, 1H), 1.1 (d, 3H, $J = 7.36$ Hz). ¹³CNMR: 166.9, 133.5, 133.4, 131.3, 130.3, 129.2, 127.7, 122.9, 122.3, 74.1, 34.8, 17.6. Anal. Calcd for C₁₄H₁₄O₂: C, 78.48; H, 6.59. Found: C, 78.35; H, 6.53.

(1S, 6S)-6-Ethyl-2,4-cyclohexadien-1-ol (4a): Following the typical procedure Cu(OTf)₂ (11.6 mg, 0.03 mmol), (*R,R,R*)-**2** (32.4 mg, 0.06 mmol), arene oxide **1a** (188 mg, 2.0 mmol), Et₂Zn (2.7 ml), 1h at -78°C (95% conversion), afforded after the usual work-up a crude reaction mixture (248 mg) constituted by the regioisomers **4a** and **6a** in a 38:62 ratio and the chiral ligand (*R,R,R*)-**2**. After repeated chromatographic purifications (SiO₂, hexanes/Et₂O=85:15 plus 1% NEt₃) it was possible to obtain an analytical pure sample of **4a** (15 mg), a colorless liquid; ¹H NMR δ (CDCl₃): 5.97-5.71 (m, 4H), 4.03-3.94 (m, 1H), 2.35-2.21 (m, 1H), 1.46-1.28 (m, 2H), 0.89 (t, 3H, $J = 7.4$ Hz). ¹³C NMR δ (CDCl₃): 132.2, 131.5, 127.6, 125.4, 69.1, 44.6, 25.4, 12.0. It was not possible to isolate in a pure state the achiral γ -adduct **6a**. However the presence of **6a** as a reaction product was clearly indicated by ¹H and ¹³C NMR of the mixture. **6a**: ¹H NMR δ (CDCl₃): 5.97-5.71 (m, 4H), 4.47-4.38 (m, 1H), 2.65-2.52 (m, 1H), 1.46-1.28 (m, 2H), 0.77 (t, 3H, $J = 7.4$ Hz). The enantiomeric excess of **4a** (64%)

was determined by chiral GC (CP-cyclodex- β -column), isothermal 107 °C, (1*S*, 6*R*) (minor) t_R 22.3 min, (1*R*, 6*S*) (major) t_R 22.8 min.

Bicyclo [4.3.0]-2-methyl-nona-3,5-dien-1-ol (3b): Following the typical procedure Cu(OTf)₂ (13 mg, 0.034 mmol), (*R,R,R*)-**2** (36.3 mg, 0.067 mmol), arene oxide **1b** (300 mg, 2.24 mmol), Me₂Zn (1.7 ml), 3h at -78°C (ca. 95% conversion), afforded after the usual work-up a crude reaction mixture (280 mg) constituted by the regioisomer **3b** and **5b** in a 80:20 ratio and the chiral ligand (*R,R,R*)-**2**. Chromatographic purification (SiO₂, hexanes/Et₂O=80:20 plus 1% NEt₃) afforded pure **3b** (80 mg, 24%) as a colourless liquid. R_f =0.38 (hexanes/Et₂O=8:2); $[\alpha]_D^{23}$ = -427.7 (c =1.77, MeOH). ¹H NMR δ (C₆D₆): 5.78 (dd, 1H, J = 9.3 and 5.2 Hz), 5.62-5.49 (m, 2H), 2.62-2.40 (m, 1H), 2.11-1.75 (m, 4H), 1.59-1.43 (m, 2H), 0.71 (d, 3H, J = 7.3 Hz). ¹³C NMR δ (C₆D₆): 147.0, 131.0, 122.5, 116.0, 79.4, 40.3, 35.7, 30.5, 23.2, 15.5.

The enantiomeric excess of **3b** (\geq 95%, not baseline separation) was determined at 220 nm by chiral HPLC (Daicel Chiralcel OD-H column), hexanes / 2-propanol 99:1, flow rate 0.5 mL/min, (-) t_R 10.9, (+) t_R 11.5 min.

Following the typical procedure Cu(OTf)₂ (13 mg, 0.034 mmol), (*R,R,R*)-**2** (36.3 mg, 0.067 mmol), arene oxide **1b** (300 mg, 2.24 mmol), Et₂Zn (3.1 ml), 3h at -78°C (95% conversion), afforded after the usual work-up a crude reaction mixture (320 mg) constituted by the regioisomer **4b** and **6b** in a 78:22 ratio and the chiral ligand (*R,R,R*)-**2**. After repeated chromatographic purifications (SiO₂, hexanes/Et₂O=85:15 plus 1% NEt₃) it was possible to obtain pure **4b** (36 mg) and **6b** (10 mg).

Bicyclo [4.3.0]-2-ethyl-nona-3,5-dien-1-ol (4b): a semisolid; R_f =0.33 (hexanes/Et₂O=85:15); $[\alpha]_D^{23}$ = -178.2 (c =0.78, MeOH). ¹H NMR δ (C₆D₆): 5.84 (dd, 1H, J = 9.7 and 5.4 Hz), 5.68-5.54 (m, 2H), 2.51-2.37 (m, 1H), 2.13-1.74 (m, 4H),

1.62-1.44 (m, 2H), 1.42-1.19 (m, 2H), 0.81 (t, 3H, $J = 7.2$ Hz). ^{13}C NMR δ (C_6D_6): 148.3, 129.1, 123.9, 116.4, 79.3, 46.7, 35.9, 30.5, 24.4, 23.4, 12.2.

Bicyclo [4.3.0]-4-ethyl-nona-2,5-dien-1-ol (6b): a liquid; $R_f=0.38$ (hexanes/ $\text{Et}_2\text{O}=85:15$); ^1H NMR δ (C_6D_6): 5.96-5.83 (m, 1H), 5.68-5.59 (m, 1H), 5.46 (d, 1H, $J = 9.5$ Hz), 2.78-2.40 (m, 2H), 2.25-1.96 (m, 4H), 1.87-1.30 (m, 3H), 0.98 (t, 3H, $J = 7.4$ Hz). ^{13}C NMR δ (C_6D_6): 149.9, 125.1, 116.9, 78.4, 48.5, 40.2, 30.8, 30.1, 23.4, 21.8, 13.0.

Regiodivergent kinetic resolution of naphthalene oxide (8).

Following the typical procedure $\text{Cu}(\text{OTf})_2$ (5.8 mg, 0.015 mmol), (*R,R,R*)-**2** (16.2 mg, 0.03 mmol), arene oxide **8** (144 mg, 1.0 mmol), Et_2Zn (1.4 ml), 3h from -78°C up to -10°C (>98% conversion), afforded after the usual work-up a crude reaction mixture (190 mg) constituted by the regioisomer (1*S*, 4*S*)-**9** and (1*R*, 2*S*)-**10** in a 66:34 ratio (as determined by ^1H NMR analysis of the crude reaction mixture) and the chiral ligand (*R,R,R*)-**2**. Dihydronaphthols **9** and **10** are not separable by chromatography on SiO_2 . However it is possible to obtain racemic **9** (isomer **10** < 5%) following the typical procedure employing $\text{Cu}(\text{OTf})_2$ (5.8 mg, 0.015 mmol), (*R,R,R*)(*S,S,S*)-**2** (16.2 mg, 0.03 mmol), arene oxide **8** (144 mg, 1.0 mmol), Et_2Zn (1.4 ml), 3h from -78°C up to -10°C . The usual work-up afforded a crude reaction mixture (160 mg) containing **9** and **10** in a 97:3 ratio. After chromatographic purification (SiO_2 , hexanes/ $\text{Et}_2\text{O}=80:20$ plus 1% NEt_3) compound **9** (90 mg, 52%) (contaminated with ca. 6% isomer **10**) was obtained.

(±)-4-Ethyl-1,4-dihydronaphth-1-ol (9): ^1H NMR δ (CDCl_3): 7.69-7.64 (m, 1H), 7.35-7.09 (m, 3H), 6.10-5.92 (m, 2H), 5.04-4.96 (m, 1H), 3.20-3.12 (m, 1H), 1.85-1.63 (m, 2H), 0.62 (t, 3H, $J=7.3$ Hz). ^{13}C NMR δ (CDCl_3): 138.62, 138.27, 131.14, 130.24, 128.11, 127.91, 127.04, 65.62, 40.87, 31.16, 10.46.

All the attempted analyses of the enantiopurity of alcohol **9** both by HPLC- and GC-CSPs gave extensive decomposition of the compound.

(+)-(1*R*, 2*S*)-2-Ethyl-1,2-dihydronaph-1-ol (10): ^1H NMR δ (CDCl_3): 7.39-7.11 (m, 4H), 6.51 (d, 1H, $J=10.0$ Hz), 6.01 (dd, 1H, $J=9.5$ and 4.6 Hz), 4.40 (d, 1H, $J=4.7$ Hz), 2.53-2.47 (m, 1H), 1.59-1.26 (m, 2H), 0.97 (t, 3H, $J=7.3$ Hz). ^{13}C NMR δ (CDCl_3): 136.44, 133.04, 131.42, 129.13, 128.38, 128.29, 127.14, 126.75, 72.77, 44.72, 25.18, 12.20. The enantiomeric excess of **10** (>98%) was determined at 254 nm by chiral HPLC (Daicel Chiralcel OD-H column), hexanes / 2-propanol 96:4, flow rate 0.5 mL/min, (-) t_{R} 15.0, (+) t_{R} 16.5 min.

The absolute configuration of compound (1*R*, 2*S*)-**10** was demonstrated by a single crystal X-ray analysis after derivatization of the enantiomer (1*S*, 2*R*)-**10** with a chiral auxiliary derived from 4,5-dichlorophthalic acid and (1*S*, 2*R*, 4*R*)-(-)-2,10-camphorsultam.²

² N. Harada, N. Koumura, M. Robillard. *Enantiomer* 1997, **2**, 303